

KOMAN, Andrej, tehn. (Zagreb)

Technical consultation of Yugoslav textile workers on the use of polyester fibers in the textile industry. Nova proizvod 15 no.5:385-387 0 '64.

Consultation on the productivity and quality of Yugoslav spinning and weaving mills. Nova proizvod 15 no.5:387-388 0 '64.

KOMAN, Andrej

~~_____~~
The second meeting of cotton spinners. Nova proizv no.3/4:236-
240 S '63.

KOMAN, Andrej

Conference on chemical fibers in the textile industry of
Yugoslavia. Kem ind 10 no.3:92 Mr '61.

FARKAS, Eve; KOMAN, Elisabeth

Artificial hibernation and infectious diseases. Acta paediat. acad.
sci. Hung. 2 no.2:99-108 '61.

1. Service de pediatrie No.III de l'Hôpital Lazlo, Budapest.
(HIBERNATION, ARTIFICIAL) (COMMUNICABLE DISEASES therapy)

FARKAS, Eva; KOMAN, Elisabeth

Electrocardiographic studies in the postinfectious stage of penicillin-treated scarlet fever. Acta pediat. acad. sci. hung. 3 no.4:381-388 '62.

1. Second Paediatric Department (Head: Dr. E. Farkas), Laszlo Hospital, Budapest (Director: Dr. J. Roman).

(ELECTROCARDIOGRAPHY) (PENICILLIN)
(SCARLET FEVER)

FARKAS, Eva, dr.; KOMAN, Erzsébet, dr.

Hibernation and infectious diseases. Orv.hetil. 102 no.31:1464-1467
30 J. '61.

1. Budapest Fovarosai Laszlo Korhaz III. sz. Gyermekosztalya.

(HIBERNATION ARTIFICIAL) (COMMUNICABLE DISEASES ther)

KARPINISHAN, K.; KOMAN, K.; KONSTANTINESKU, K.; BADIYA, D.

Significance of a mechanical suture in preventing bronchial
fistulae following lung resections. Grud. khir. 6 no.1.76-78
Ja-F '64. (MIRA 18:11)

1. Klinika grudnoy khirurgii (sav. - prof. K. Karpinishan)
bol'nitsy "Filaret", Bukharest. Adres avtorov: Bukharest,
klinika grudnoy khirurgii bol'nitsy "Filaret". Submitted
March 25, 1963.

KARPINISHAN, K. [Carpinisan, C], prof., doktor; BOGDAN, Tr. [Bogdan, Traian],
kand.med.nauk, doktor; KOMAN, K. [Coman, C], doktor

Decortication of the lung. Vest. khir. 90 no.3:30-35 Mr'63.
(MIRA 16:10)

1. Iz Bukharestskoy kliniki grudnoy khirurgii (dir. - prof.
doktor K.Karpinishan).

(LUNGS—DISEASES)

(PLEURA—SURGERY)

ROMAN, M.

Definition of the Topological K -Linear ¹⁶

Roman, Milan. Bemerkung zu einer Definition der topologischen K -Lineale. Časopis Pěst. Mat. 83 (1958), 155-159. (Czech. Russian and German summaries)

J. Matík [Časopis Pěst. Mat. 79 (1954), 3-40; MR 16, 492] has defined a normed K -lineal (vector lattice) as a K -lineal in which $\|a\| = \|a\|$ and $0 \leq a \leq b$ implies $\|a\| \leq \|b\|$. Let H be the linear space of all real functions on $[0, 1]$ of the form $f+g$, where f is continuous and g vanishes except on a finite set. Let H have the usual linear operations and order, and set $\|f+g\| = \max |f(x)| + \sum_{0 \leq x \leq 1} |g(x)|$. Then H is a normed K -lineal in which the algebraic and order operations are continuous, but $0 \leq a \leq b$ does not imply $\|a\| \leq \|b\|$. In fact, there is a neighborhood U of 0 such that for every neighborhood V of 0 there are elements a, b for which $0 \leq a \leq b$, $b \in V$, and $a \notin U$. Furthermore, the linear functional $J(f+g) = \sum g(x)$ is continuous, but its positive part J_+ is infinite at 1 ($J_+(1) = \sup J(h)$; $0 \leq h \leq 1$).

E. Hewitt (Seattle, Wash.)

2
1-FW

25069

S/080/60/033/010/023/029
D216/D306

158540

AUTHORS: Mikhail, R., Alexsandru, L., Koman, M., and
Yurchenko, V.

TITLE: Modified polyethylene terephthalate as an
electro-insulating varnish

PERIODICAL: Zhurnal prikladnoy khimii, v. 33, no. 10, 1960,
2336 - 2340

TEXT: To obtain modified polyethylene terephthalate three routes were taken: 1) Introduction of all components into general reaction; 2) Transesterification of dimethyl terephthalate (DMT) with corresponding glycols, followed by polycondensation (all reagents entering general reaction); 3) Products from transesterification for the given glycol separated and then polycondensed. The basic study was done on polyethylene terephthalate modified with glycerol, i.e. the effect of change in (a) molar proportions of components (b) temperature of the reaction and (c) time of reactions.

Card 1/4

25069

S/080/60/033/010/023/029

D216/D306

Modified polyethylene ...

tion, 30 moles of dimethyl terephthalate were used, 50, 60, 65 moles of ethylene glycol and 20, 10, 5 moles of pentaerythrite. On polycondensation of dimethyl terephthalate, ethylene glycol, glycerol and pentaerythrite in proportions 25:50:22:3 respectively and at 190°C for 180 min and at 3 mm Hg, a soluble transparent product was obtained with a melting point of 85° and 350 OH groups. On polycondensation of dimethyl terephthalate, ethylene glycol and glycerol with proportions 40:40:20 at 240°C for 270 min., a transparent soluble product is obtained with a melting point of 95°C and 377 OH groups. Synthesized products had molecular weights from 1200 to 1400 and these were determined by the cryoscopic method, in phenol. Use of these varnishes on copper conductors has given resistance to 5000 volts potential, thermal stability up to 155°C, and good resistance to wear. Especially good results were obtained with the varnish based on polyethylene terephthalate modified with ethylene glycol, glycerol, pentaerythrite. There are 6 figures, 1 table and 2 references: 1 Soviet-bloc and 1 non-Soviet-bloc. X

Card 3/4

25069

Modified polyethylene ...

S/080/60/033/010/023/029
D216/D306

ASSOCIATION: Nauchno-issledovatel'skiy khimicheskiy institut
Bukharest (Scientific-Research Chemical Institute,
Bucharest)

SUBMITTED: February 19, 1960

Card 4/4

ALEKSANDRU, L. [Alexandru, L.]; KOMAN, M. [Coman, M.]; RIZESKU, T.
[Rizescu, T.]; POPOVICH, A. [Popovici, A.]

Effect of the nature of stabilizers on the stability of polycaprolactam. Khim.volok no.4:4-8 '62. (MIRA 15:8)

1. Nauchno-issledovatel'skiy institut khimii, Bukharest, Rumynskaya Narodnaya Respublika.

(Nylon)

KOMAN, Milan (Praha); ROHLICEK, Jiri (Praha)

"Mathematical methods in economic practice" by Anton Kolzig.
Reviewed by Milan Koman and Jiri Rohlicek. Cas pro pest mat
88 no.2:254-256 '63.

//, sept. components have sharp borders.

Jan Mlicka

Koman V

COUNTRY : Czechoslovakia
CATEGORY :

ABS. JOUR. : RZKhim., No. 21 1959, No. 74696

AUTHOR : Palo, V., Koman, V., and Hrabe, Z.
INST. : Not given

TITLE : The Separation of Higher Fatty Acids by Paper Chromatography by Varying the Concentration of Solvent During Elution

ORIG. PUB. : Chem Zvesti, 12, No 9, 525-532 (1958)

ABSTRACT : The authors propose the separation of the higher fatty acids by paper chromatography with continuous variation of the concentration of the mobile phase. A 50-200 γ sample of a mixture of fatty acids (FA) is deposited in the form of a 5% solution in C_6H_6 on Whatmann No 3 paper impregnated with a 10% solution of paraffin oil (sp gr 0.880-0.895) in C_6H_6 . The paper is placed between glass plates and chromatographed for 12 hrs at 20° with 50% CH_3COOH ;

CARD: 1/3

86

COUNTRY : Czechoslovakia
CATEGORY :

ABS. JOUR. : RZKhim., No. 21 1959, No. 74696

AUTHOR :
INST. :
TITLE :

ORIG. PUB. :

ABSTRACT : the concentration of the latter is gradually increased during the run. The chromatograms are dried at 120°, dipped for 45 min in a solution of Cu acetate (20 ml of saturated solution + 240 ml water); this converts the FA spots to the corresponding Cu salts. The excess Cu acetate is rinsed off with water and the chromatogram is treated with 7.5% $K_4Fe(CN)_6$ which gives the spots a red-brown color. The following R_f values were obtained for the acids listed: erucic 0.02, stearic 0.02, palmitic 0.12, oleic 0.17, myristic

CARD: 2/3

KOMAN, Vaclav

Changes in rape oil fat acids during refining and hydrogenation.
Prum petravin 15 no.8:394-395 Ag '64.

1. Slovak Higher School of Technology, Faculty of Chemistry,
Bratislava.

MICHALEC, U.; SULC, M.; MESTAN, J.; HALAMA, D.; KOMAN, V.

Lipids in certain pathological conditions. II. Studies on fractions on cholesterol esters in human and animal blood. Sborn. lek. 63 no.4:99-103 Apr 1961.

1. Laborator pro metabolismus bikovin a proteosyntezu fakulty vseobecneho lekarstvi University Karlovy v Praze, reditel prof. dr. J.Horejsi, Laborator angiologicka fakulty vseobecneho lekarstvi University Karlovy v Praze, reditel prof. dr. B.Prusik, Slovenska vysoka skola technicka, katedra biochemie a mikrobiologie v Bratislave, reditel prof. dr J.Nemec. (CHOLESTEROL blood)

KOMAN, Vachy

~~Dr. Vachy Komarov~~

Dr. Vachy Komarov

Dr. Vachy Komarov

Affiliations:

Source: Bratislava, Dr. Vachy Komarov, No 10, Oct 60, p 600

~~Dr. Vachy Komarov~~

Source: Engineer
Affiliations: Department of Technical Microbiology and Biochemistry
at the Slovak Technical University in Bratislava.

Data: Co-author of "Quantitative Determination and Analysis of
Free-chloric Acid during Hardening of Sunflower-seed Oil by
Means of Paper Chromatography," Source.

Source: Engineer
Affiliations: Department of Technical Microbiology and Biochemistry
at the Slovak Technical University in Bratislava.

Data: Co-author of "Quantitative Determination and Analysis of
Free-chloric Acid during Hardening of Sunflower-seed Oil by
Means of Paper Chromatography," Source.

[Handwritten signature]

KOMANAR, S. I.

PHASE I BOOK EXPLOITATION SOV/5289
Akademiyu nauk SSSR. Tekhnicheskaya nauchno-issledovatel'skaya laboratoriya elektricheskoy obrabotki materialov.
Elektroiskrovaya obrabotka metallov (Electric-Spark Machining of Metals) no. 2. Moscow, Izd-vo AN SSSR, 1950. 262 p. Errata slip inserted. (Series: Its: Trudy) 6,000 copies printed.
Sponsoring Agency: Akademiyu nauk SSSR.

Resp. Ed.: B. R. Lazarenko; Ed. of Publishing House: S. M. Mozhes; Tech. Ed.: A. P. Uuseva.

PURPOSE: This collection of articles is intended for process engineers, and technical and research personnel engaged in the working of metals.

COVERAGE: Problems concerning the most effective application of electric-spark methods in industry are reviewed. Possible future developments in the field of electric-spark machining and its automation are discussed, and, for instance of its present utilization in industry, the technical-economic effectiveness of the process is examined, and the equipment involved is described. The relationship between the parameters of electric-spark pulse accuracy, and the production characteristics (productivity, machining accuracy, and surface quality) of electric-spark machining is established. An electric-spark method of advanced cutting of wire, thus directly producing a finished part, 30 micron thick, is described. The use of electric-spark machining for the treatment of metal surfaces is also mentioned. There are 121 references: 82 Soviet, 20 English, 10 French, 8 German, and 1 Italian. These references accompany individual articles.

Zolotykh, B. N., and I. P. Korobova. Selecting Optimum Regimes for Electric-Spark Machining of Sintered-Carbide Alloys 114
Chetverikov, S. S., and N. K. Poteyev. Electric-Spark Machining of the Cutting Elements of High-Carbon-Alloy Blanking Punch-Die Sets 120
Gul'yan, K. E. The Electric-Spark Method Applied to Threading 142
Kholodnov, Ye. V. Manufacture of Precision Tools by the Electric-Spark Method 155
Gul'yan, K. E., and V. L. Kravchenko. Manufacture of Complex-Shaped Machine Parts by Using a Program-Controlled Electric-Spark Machining Unit 179
Aleksandrov, V. P., and B. N. Zolotykh. Selecting the Optimum Procedures for Electric-Spark Machining of Nickel-Base Heat-Resistant Alloys 196
Gorbunov, B. N. Electric-Spark Lapping Used on Flour-Mill Rolls 205
Fronko, G. P. Manufacture of Stainless and High-Manganese Steel Parts by the Electric-Spark Method 217
Ayzenshtok, V. L., and S. I. Komanar. Electric-Spark Machining of Mass-Produced Parts 227
Levinson, Ye. M. The Development of Electric-Spark Machining in Mass Production 233
Card 4/5

KOMAN, Vaclav, inz. (Bratislava, Kollarovo namesti 2, Chemicky pavilon, Slovenska vysoka skola technicka); KOVAC, Stefan, dr., inz., C.S. (Bratislava, Kollarovo namesti 2, Chemicky pavilon, Slovenska vysoka skola technicka); KOMANOVA, Eva, inz.

Infrared absorption spectra of esters of the trans - Δ^9 -octadecenoic acid (elaidic acid). Chem zvesti 15 no.6:441-449 Js '61.

1. Katedra technickej mikrobiologie a biochemie, Slovenska vysoka skola technicka, Bratislava (for Koman).
2. Katedra organickej chemie, Slovenska vysoka skola technicka, Bratislava (for Kovac).
3. Vyvojove pracovisko, n.p., Palma, Bratislava, ul. Februaroveho vitazstva (for Komanova).

KOVAC, Stefan, dr., inz., C.Sc. (Bratislava, Kollarovo namesti 2, Chemicky pavilon, Slovenska vysoka skola technicka); KOMAN, Vaclav, inz. (Bratislava, Kollarovo namesti 2, Chemicky pavilon, Slovenska vysoka skola technicka); KOMANOVA, Eva, inz.

Examination of the effect of symmetry of esters of the trans- Δ^9 -octadecenoic acid (elaidic acid) on the altitude of infrared absorption band in 970 cm^{-1} . Chem zvesti 15 no.6:450-455 Je '61.

1. Katedra organickej chemie, Slovenska vysoka skola technicka, Bratislava (for Kovac). 2. Katedra technickej mikrobiologie a biochemie, Slovenska vysoka skola technicka, Bratislava (for Koman). 3. Vyvojove pracovisko, n.p., Palma, Bratislava, ul. Februaroveho vitazstva (for Komanova).

FEDOROVA, N. Ye.; MORYGANOV, P. V.; ROMANENKOVA, L. A.

Mechanism of the action of the stabilizers of hydrogen
peroxide alkali solutions and its practical application.
Izv. vys. ucheb. zav.; tekhn. tekst. prom. no. 4:76-83 '62.
(MIRA 15:10)

1. Ivanovskiy khimiko-tekhnologicheskii institut.

(Bleaching) (Hydrogen peroxide)

MATOV, Konstantin; KOMANDAREV, Stamat

The intensity of invasion of *Trichinella* in various swine muscles.
Wiad. parazyt. 8 no.6:613-628 '62.

1. Kafedra parazitologii Vet. Insti, Sofiya, Bulgariya.
(TRICHINOSIS) (SWINE DISEASES)

MATOV, K.; KOMANDAREV, St.

Further studies on the problems of muscular Trichinella
larvae occurring in the organs having no striated muscles.
Izv khelminth lab BAN 9:81-90 '64.

KOMANDAREV, St.; IOVCHEV, E.

Studies on the mechanism and factors of immunity in
trichinellosis. Izv khelmin lab BAN 9:117-123 '64.

TOMOV, Vl.; KOMANDAREV, St.

Possibility of preventing recurrences of echinococcosis in man
by applying the proteolytic enzyme Papain. Izv khelminth lab
BAN 9:129-134 '64.

KOMANDENKO, N.I.

Effect of oxygen under pressure on the course of experimental tick-borne encephalitis in white mice. Vop.virus. 7 no.6:658-661 N-D '62. (MIRA 16:4)

1. Voenno-meditsinskaya ordena Lenina akademiya imeni S.M. Kirova, Leningrad.

(ENCEPHALITIS)

(OXYGEN THERAPY)

KOMANDENKO, N.I.

Treatment of the progressive form of Russian tick-borne
encephalitis with oxygen under pressure. Vop. psikh. i nevr.
no.9:250-256 '62. (MIRA 17:1)

1. Kafedra nervnykh bolezney Voenno-meditsinskoy ordena
Lenina akademii imeni S.M. Kirova.

KOMANDI, Gyorgy

Four-wheeled tractors. Jarmu mezo gep 7 no.8:303-306 '60.

DEMICHEV, A.D.; KISELEV, V.F., starshiy dorozhnyy master (stantsiya Ira-Iol' Pechorskoy dorogi); KOZLOVSKIY, A.D.; KOMANDIN, A.A., starshiy dorozhnyy master (stantsiya Polotsk Belorusskoy dorogi); KURS, V.G., brigadir puti (stantsiya Cherekhovo Vostochno-Sibirskoy dorogi); PAVLOV, V.N., brigadir puti (stantsiya Cherekhovo Vostochno-Sibirskoy dorogi); SHAKHBALAYEV, A.M., dorozhnyy master (stantsiya Zenzeli Ordzhonikidzevskoy dorogi); TARASENKO, V.Ye., dorozhnyy master (stantsiya Irkutsk II)

Letters to the editor. Put' i put.khoz. no.11:43-45 N '58.

(MIRA 11:12)

1. Nachal'nik normativnoy stantsii tresta "Rekput'." (for Demichev).
2. Zamestitel' nachal'nika distantsii, stantsiya Kizel Sverdlovskoy dorogi (for Kozlovskiy).

(Railroad engineering)

KOMANDIN, Arnol'd Grigor'yevich; SANOYLOVICH, T.A., red.

[Operation of a MAK-FV-4 freon refrigeration plant]
Ekspluatatsiia freonoi kholodil'noi ustanovki marki
MAK-FV-4. Moskva, Izd-vo "Transport," 1964. 51 p.
(MIRA 17:6)

PROCEDURES AND PROPERTIES INDEX

82

Influence of an inert solid phase on the f.p. of water and dilute aqueous solutions. III.

Amount of freezable water in presence of quartz sand and starch. D. N. TARASENKOV and A. V. KOMAROV (J. Gen. Chem. Russ., 1936, 6, 1147-1160).—The proportion of H₂O not freezing at 0° or total surface of the SiO₂ particles in unit vol., or to the concn. of starch, and may amount to 33% in the former, and to 70% in the latter, case. R. T.

ASM-A14 METALLURGICAL LITERATURE CLASSIFICATION

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

TARASENKOV, D. N.; KOMANDIN, A. V.

"Vapor Tension of Pentachlorides of Tantalum and Niobium and the Mixtures with Tetrachloride of Titanium," Zhur. Obshch. Khim., 10, No. 14, 1940. Lab. of Chem. Thermodynamics. Moscow State Univ. and State Sci. Res. Inst. of Rare Elements. Received 25 March 1940.

Report U-1610, 3 Jan. 1952.

147 AND 148 PAGES

PROCESSES AND SUBSTITUTED MODELS

Physicochemical investigation of tungsten oxychlorides.
A. V. Komandir and D. N. Tarasenkova. J. Gen. Chem. (U. S. S. R.) 15, 1535-42 (1940). The various methods for prepn. of WO_2Cl_2 were compared. A max. yield of most pure WO_2Cl_2 prepd. by the chlorination of WO_3 with the Cl-N mist. in ratio 2:3, was obtained at 500 $^{\circ}C$. The vapor pressure of WO_2Cl_2 detd. in the Cl stream at 165-204 $^{\circ}C$ varied from 0.10 to 3.00 mm. of Hg. The dissociation of WO_2Cl_2 , according to the equation $2WO_2Cl_2 \rightleftharpoons WOCl_4 + WO_3$, started at 250 $^{\circ}C$. The total pressure at 268-434 $^{\circ}C$ varied from 2.40 to 474.0 mm. and at the same temp. the partial pressure of $WOCl_4$ varied from 1.12 to 100.0 mm. The vapor pressure of the WO_2Cl_2 + 30% $WOCl_4$ and 20% WO_2Cl_2 + 80% $WOCl_4$ mists, at 164-265 $^{\circ}C$ and 165-178 $^{\circ}C$, resp., varied from 48.0 to 67.9 mm. and from 18.0 to 46.8 mm. The solubilities of WO_2Cl_2 and $WOCl_4$ in $PhCl$ at 25 $^{\circ}C$ were 0.15% by wt. and at 114 $^{\circ}C$ was 0.29 and 0.43%, resp. The vapor phase of the solid solns. of W oxychlorides in $PhCl$ at their h. p. contained no WO_2Cl_4 and only traces of $WOCl_4$. The chlorination of the W + WO₃ mist. at 300-500 $^{\circ}C$ yielded 87-89% of WO_2Cl_2 , the vapor pressure of which, measured in the Cl stream, varied at 100-50 $^{\circ}C$ between 1.8 and 9.4 mm. and at 170-214.5 $^{\circ}C$ between 188 and 701 mm. of Hg. A. A. Podernov

State Sci. Res. Inst. Rare Elements
Lab. Chem. Thermodynamics, Moscow State U.

ASB-15A METALLURGICAL LITERATURE CLASSIFICATION

FROM SYNDICATE

RECORDS - 119 NOV 1941

RECORDS - 119 NOV 1941

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

Technology

Electromotive forces. Moskva, Izd-vo Moskovskogo universiteta, 1951. Pod red. A. V. Frosta, Moskva.

9. Monthly List of Russian Accessions, Library of Congress, June 195~~8~~⁷, Uncl.

2

CA

Undercooling of organic liquids. Effect of various factors on the undercooling of benzene. A. N. Kozmandin (Moscow State Univ.). *Technical Mater. Sci.* 8, No. 10, 1967, 1161-1162. (English transl. No. 6, 55-56 (1967)). Benzene, purified chemically, and by fractional freezing and distn., to a sharp m.p. of $5.43 \pm 0.02^\circ$, and outgassed, could be undercooled to 4.5° below the freezing temp. in vessels of 10 mm. or larger diam., and kept liquid for 25 hrs. provided the liquid was not mechanically disturbed, e.g. by shaking. Stirring of liquid C_6H_6 undercooled to only 0.5° produced immediate crystals. C_6H_6 with air does not counteract undercooling, as air-satd. C_6H_6 could be undercooled by about the same amt. as outgassed C_6H_6 . Agitated C_6H_6 , by about the same amt., cannot be undercooled by more than $0.1-0.2^\circ$; the material of the agitator (metal or glass) makes no difference. In capillaries of 0.2 mm. diam., C_6H_6 can be undercooled by about 70° . A strong jump of the max. possible undercooling occurs between a vessel of 3.0 and one of 2.33 mm. diam.; in the former, the undercooling is still 4.5° , whereas in the next narrower tube it is possible to undercool by 10.5° , and in a 0.44-mm. tube, by 20.1° . N. Thon

KOMANDIN, A. V.

Sep 52

USSR/Chemistry Vanadium Compounds

"Vapor Pressure of Vanadium Oxytrichloride," A. V. Komandin and M. L. Vlodavets, Moscow State U

Zhur Fiz Khim, Vol 26, No 9, pp 1291-1297

Made a comparative study and evaluation of two methods of obtaining VOCl_3 . The method of chlorinating V_2O_3 with Cl_2 gas in the presence of C proved to be more effective than the action of dry HCl gas on V_2O_5 . By the first method, the chlorination of 30 grams of V_2O_3 proceeded quantitatively and to completion in 2 hrs. The vapor pressure of VOCl_3 was measured within the temp range of $18.7-100^\circ$. The heat of evapn and the Trouton Kistyakovskiy const were computed from the vapor pressure temp relationship. From the heating curves, the MP of VOCl_3 was found to be $-78.9 \pm 0.2^\circ$.

263 T 19

REMANDED, A-1

Supercooling of organic liquids
 George Joseph Bristow, 1952-1953, 1954-1955
 Dissertation No. 104, 124-224 (1953). - Supercooling of
 and purified glycerol (I), triethylamine (II), tri-
 phenylene (III), benzene (IV), and acrylonitrile and
 was studied under various conditions. In water con-
 tainers IV could be supercooled no more than 9° below its
 m.p.; however in capillary tubes (about 0.2 mm in diam)
 it was supercooled to 70° below the m.p. and was kept in
 this condition for about 10 hrs. In glass tubes of 1 mm
 diam. and 100 mm in length it was possible to supercool I
 by 22.5°. A few samples of I and III have been kept in the
 supercooled state for 6 yrs. at 10-30°. When I was cooled
 in 100° in sealed tubes for 10 min. and then cooled to
 room temp. it remained a clear viscous liquid for 3 yrs.
 was impossible to supercool II under any conditions.
 Addition of water to I causes very rapid crystal-
 lization. The rate of crystal growth increased as the concn. of water was increased from
 1.5% to 40%, at which point it ceased to increase. Categori-
 zation of liquids into 3 categories with respect to super-
 cooling is suggested: (1) liquids of the benzene type which
 type composed of small molecules and molecules which are
 molecules with considerable translational motion. These liquids
 undergo crystal- very easily and are difficult to supercool.
 (2) liquids of the molal type with relative large molecules
 less sym. than those in the 1st class, and molecules with
 rotational brownian motion. Coordination of these liquids
 liquid state is similar to that in the crystal state. These
 liquids are very easily supercooled. (3) high polymers and
 polymeric liquids which are rather large molecules and
 usually long, branched, more or less irregular in
 shape and within the same mass are very difficult to
 supercool. These liquids are very difficult to supercool and
 crystallize and are in their usual state supercooled liquids.

KOMANDIN, A.V.

Use of a chlorine-Ag electrode instead of a calomel electrode.
Uch.zap.Mosk.un. no.164:201-207 '53. (MIRA 8:7)
(Electrodes, Silver)

KOMANDIN, A. V.

USSR/ Chemistry Physical chemistry

Card : 1/1

Authors : Komandin, A. V., and Bonetskaya, A. K.

Title : Dipole moments of certain salicylic acid derivatives

Periodical : Zhur. fiz. khim. 28, No. 6, 1113 - 1119, June 1954

Abstract : The dipole moments of methyl salicylate, ethyl salicylate, isoamyl salicylate, phenyl salicylate and 3-naphthyl salicylate, were measured in benzene solutions and the dipole moments of ortho-acetoxybenzoic acid in benzene solution at 25°C. The effect of the change in dipole moments in many salicylic acid derivatives and the effect of the intra-molecular hydrogen bond on the change of the dipole moments are explained. Drawing of the device used in measuring the dielectric permeability of the solutions, is included. Twenty references. USSR, 16 German and English. Tables.

Institution : The M. V. Lomonosov State University, Moscow

Submitted : November 13, 1953

AY

147-12/25

Authors : Komandin, A. V., and Bonetskaya, A. K.

Title : Dipole moments of orthohydroxybenzoic acid substitutes

Periodical : Zhur. fiz. khim. 28/10, 1789-1794, Oct 1954

Abstract : In order to explain the effect of the hydrogen bond on the dipole moment, the authors measured the dipole moments of the following substitutes of orthohydroxybenzoic acid: orthomethoxy- and orthobenzoxybenzoic acid, methyl ether of orthomethoxybenzoic acid and salicylic acid (the latter represents a molecular compound of salicylic acid and acetylene). Data regarding the measured dielectric permittivity, density and index of refraction for various concentrations of the dissolved substances are presented. The chemical structure of the substitutes is described. Twelve references: 1-USSR; 2-German; 2-USA and 1-French (1893-1954). Tables.

Institution : The M. V. Lomonosov State University, Moscow

Submitted : February 17, 1954

USSR/Chemistry - Physical chemistry

Card 1/2 : Pub. 147 - 18/27

Authors : Komandim, A. V., and Rosolovskiy, V. Ya.

Title : Dipole moments of certain orthohydroxybenzoic acid and glycerin derivatives

Periodical : Zhur. fiz. khim. 28/12, 2215-2221, Dec 1954

Abstract : In order to explain the effect of the hydrogen bond on the dipole moment, the following compounds were measured at 25° C in benzene solutions: methyl and phenyl ethers of o-acetoxybenzoic acid (methyl and phenyl o-acetoxybenzoate), phenyl ether of o-phenoxybenzoic acid (phenyl o-phenoxybenzoate) and glycerin ether of acetic acid (glycerin acetate) as well as glycerin ether of salicylic acid (glycerin salicylate) in dioxane. The dipole moment of the latter was measured in dioxane because of its low solubility in benzene. The synthesis and purification of the investigated substances are described and the results obtained are tabulated. Fourteen references ; 2 USSR; 6 German; 1 British; 1 French; 3 USA and 1 Swiss (1897-1954). Tables.

Chem. fiz. khim. 28/12, 2215-2221, Dec 1954

(Additional Card)

Card 2/2

Institution : The M. V. Lomonosov State University, Moscow

Submitted : May 8, 1954

5(4)

SOV/76-33-3-10/41

AUTHORS: Komandin, A. V., Bonetskaya, A. K.

TITLE: Density and Molar Volume of Several Organic Compounds in a Broad Range of Temperature (Plotnosti i molyarnyye ob'yemy nekotorykh organicheskikh soyedineniy v shirokom intervale temperatur)

PERIODICAL: Zhurnal fizicheskoy khimii, 1959, Vol 33, Nr 3, pp 566 - 571 (USSR)

ABSTRACT: Density and molar volume in the temperature range 100-200° were determined for the liquid and supercooled liquid state of the following 11 substances: methyl-, ethyl-, isoamyl-, phenyl-, (salol) and 2-naphthyl- (betol)-esters of o-hydroxy benzoic acid, o-acetoxy benzoic acid (aspirin), the methyl esters of o-methoxy benzoic acid, salipyrine, o-methoxyphenol (guaiacol), 3-methyl-6-isopropylphenol (Thymol) and benzophenone. Density in solid state was determined at room temperature for: the phenyl- and 2-naphthyl esters of o-hydroxy benzoic acid, o-aceto benzoic acid, 3-methyl-6-isopropylphenol and benzophenone; these data are, however, to be re-

Card 1/2

Density and Molar Volume of Several Organic Compounds
in a Broad Range of Temperature

SOV/76-33-3-10/41

garded as orientation data only. The production of guaiacol, thymol and benzophenone is given, the way of production of the other substances has already been described (Ref 4). The density was measured by means of a pycnometer-dilatometer of the Biron-type (Ref 5). Table 2 shows that the temperature function of density develops linearly for all substances investigated. There are 3 tables and 5 references, 2 of which are Soviet.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova
(Moscow State University imeni M. V. Lomonosov)

SUBMITTED: June 25, 1957

Card 2/2

5 (4)

AUTHORS:

Komandin, A. V., Bonetskaya, A. K.
(Moscow)

SOV/76-33-5-3/33

TITLE:

The Dielectric Constant of Esters of Orthohydroxy Benzoic Acid in a Wide Temperature Interval (Dielektricheskaya pronitsayemost' slozhnykh efirov ortogidroksibenzoynoy kisloty v shirokom intervale temperatur)

PERIODICAL:

Zhurnal fizicheskoy khimii, 1959, Vol 33, Nr 5, pp 976 - 982 (USSR)

ABSTRACT:

This paper describes measurements of the dielectric constants of methyl, ethyl, isoamyl, phenyl, and 2-naphthyl ester of the acid mentioned in the title. The measurements were carried out in a liquid and undercooled liquid state within a temperature interval ranging of from a temperature below melting point to a temperature at which the dielectric constant decreased rapidly to low values with a constant frequency of the outer electric field; these values did not vary with further temperature decrease and approached the square of the refractive index, i.e. the total amount of electronic and atomic polarization. Figure 1 shows the construction of the condenser used for the measurements. By using a special glass filter crystal-

Card 1/3

The Dielectric Constant of Esters of Orthohydroxy
Benzoic Acid in a Wide Temperature Interval

SOV/76-33-5-3/33

lization germs were prevented from being carried along with the liquid and the measurement in undercooled state up to glass-shaped modification without spontaneous crystallization was rendered possible. Tables 1 - 5 show the values of the dielectric constant for the esters mentioned; moreover, the values computed for the Kirkwood coefficient g , as well as for the general and orientation polarization. Figure 2 shows the shape of the curve of the dielectric constant depending on temperature. Hence it appears that with decreasing temperature and constant frequency of the outer electric field ($1.72 \cdot 10^6$ cycles) the value of the dielectric constant increases, reaches a maximum, and then decreases rapidly to low values. There is a relation between the molecular structure of the substances investigated and the variation of the dielectric constants. The more complicated the structure of the substituent, the lower and flatter the curve of the dielectric constant. The maxima of the dielectric constants are influenced in a similar way. These turning points could be reproduced with an accuracy

Card 2/3

The Dielectric Constant of Esters of Orthohydroxy
Benzoic Acid in a Wide Temperature Interval

SOV/76-33-5-3/33

of 1 - 2° and represent a specific quantity as to the liquid
concerned. There are 2 figures, 5 tables, and 12 references,
6 of which are Soviet.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova
(Moscow State University imeni M. V. Lomonosov)

SUBMITTED: November 25, 1957

Card 3/3

5(4)

AUTHORS: Komandin, A. V., Rosolovskiy, V. Ya.

SOV/76-33-6-18/44

TITLE: Densities and Molar Volumes of Some Organic Compounds in Broad Temperature Ranges (Plotnosti i molyarnyye ob'yemy nekotorykh organicheskikh soyedineniy v shirokom intervale temperatur)

PERIODICAL: Zhurnal fizicheskoy khimii, 1959, Vol 33, Nr 6, pp 1280-1282 (USSR)

ABSTRACT: The densities were determined in the temperature range of 60 - 160° for the liquid and undercooled state of the following 6 compounds: ethylene glycol, glycerin, α, γ -glyceryl acetate, α -glycerin-o-hydrobenzoate, and the methyl- and phenyl-o-acetoxy benzoate (Table 1). The determination accuracy is specified as being $\pm 0.03\%$. Results show that in all compounds in the investigated temperature range the temperature function of density proceeds linearly. Also the molar volumes exhibit a linear temperature function. Table 2 supplies equations of the straight lines of the temperature function of density and molar volumes of the compounds investigated for broad temperature ranges. There are 2 tables

Card 1/2

SOV/76-33-6-18/44

Densities and Molar Volumes of Some Organic Compounds in Broad Temperature Ranges

and 3 references, 2 of which are Soviet.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova
(Moscow State University imeni M. V. Lomonosov)

SUBMITTED: November 10, 1957

Card 2/2

5(4)

SOV/76-33-6-19/44

AUTHORS: Komandin, A. V., Rosolovskiy, V. Ya.

TITLE: Dielectric Constant of Some Organic Compounds Over a Broad Range of Temperature (Dielektricheskaya prornitsayemost' nekotorykh organicheskikh soyedineniy v shirokom intervale temperatur)

PERIODICAL: Zhurnal fizicheskoy khimii, 1959, Vol 33, Nr 6, pp 1283-1288 (USSR)

ABSTRACT: To clarify the relationship existing between dielectric properties and structure the dielectric constants (DC) were determined for the liquid and undercooled state over a broad temperature range concerning the following substances: methyl- and phenyl-o-acetoxybenzoate, α -glyceryl-o-hydroxybenzoate and α,γ -glyceryl acetate. The preparation and cleaning methods as well as the physical constants of these compounds have already been described earlier (Ref 2). The determination of the (DC) took place according to the pulsating method (Ref 3). Measuring results of the (DC) and densities of the above mentioned substances are given as well as the computed values of the general and orientation polarization for the liquid and undercooled phase, and the

Card 1/2

SOV/76-33-6-19/44

Dielectric Constant of Some Organic Compounds Over a Broad Range of Temperature

coefficients g according to Kirkwood (Tables 1 - 4). It may be observed from the results obtained and from a graph (Figure) that the (DC) rises with the temperature drop, attains a maximum and then drops rapidly. The position of the curve maxima, however, depends on the structure of the compound. With rising molecular weight of ester the $\epsilon - T$ curve ($\epsilon = (DC)$, $T = \text{temperature}$) runs lower, and the curve maximum shifts to higher temperatures. The temperatures T_x (characteristic of each of the compounds) which corresponded to the $\epsilon - T$ curve inflection, were determined at constant frequencies of an electric outer field, and are specified in the present paper. There are 1 figure, 4 tables, and 4 Soviet references.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova
(Moscow State University imeni M. V. Lomonosov)

SUBMITTED: November 10, 1957

Card 2/2

9/076/60/034/04/24/042
B010/B009

AUTHORS: Komandin, A. V., Bonetskaya, A. K. (Moscow)

TITLE: The Dielectric Constants of Some Organic Compounds Within a Broad Temperature Range

PERIODICAL: Zhurnal fizicheskoy khimii, 1960, Vol. 34, No. 4, pp. 845 - 849

TEXT: In continuation of previous papers (Refs. 1,2) concerning the relation between the dielectric properties and chemical structures of organic substances in liquid and supercooled liquid phases the dielectric constants of the methyl esters of o-methoxybenzoic acid, o-acetoxybenzoic acid, salipyrine, and benzo-phenone were measured. The measurements were made by means of an apparatus previously described (Ref. 3) at a frequency of $1.72 \cdot 10^6$ cps and temperatures of from 200 to 400°K. The measured values of the dielectric constants and densities of the substances under investigation as well as the calculated values of the total and oriented polarizations for both the liquid and supercooled phases and of the Kirkwood coefficients g of the intermolecular interaction are given in Tables 1-4. As in the earlier investigations, the curve of the dependence of the

Card 1/2

84253

M.7800 also 2209

S/076/60/034/009/018/022
B015/B056

AUTHORS: Komandin, A. V., Shapovalova, R. D., and Mikhaylova, N. P.

TITLE: Some Physical Properties of Tungstates. II. The Dielectric Constant and the Polarization of Solid Tungstates

PERIODICAL: Zhurnal fizicheskoy khimii, 1960, Vol. 34, No. 9, pp. 2063-2065

TEXT: The dielectric constants of manganese-, calcium-, barium-, zinc-, copper-, magnesium-, iron-, cobalt-, and nickel tungstates were measured by the immersion method (Refs. 1,2) in the solid state at 25°C (Table, measured values). As standard liquids, benzene - acetone and acetone - water mixtures were used for the solid tungstates. Measurements were carried out on a previously described device (Ref. 3) at a frequency of $1.72 \cdot 10^6$ c/sec. From the values obtained for the dielectric constants, the total polarizations of the solid crystalline tungstates were calculated from the Debye equation. The dielectric constant is in the range from 17.7 to 21.4. The molar refraction for calcium- and manganese tungstate

Card 1/2

Some Physical Properties of Tungstates. II.

The Dielectric Constant and the Polarization of
Solid Tungstates.

84253

S/076/60/034/009/018/022
B015/B056

in the solid state was also determined. According to the results obtained it is found that, apparently, the structure of the crystals of all tungstates investigated is of the type of ionic crystals, and that the difference between the total polarization and the molar refraction represents the polarization of ionic displacement. There are 1 table and 6 references: 5 Soviet and 1 US.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova
(Moscow State University imeni M. V. Lomonosov)

SUBMITTED: December 31, 1958

Card 2/2

KOMANDIN, A.V.; SHIMIT, B.D.

Thermodynamics of dielectric relaxation processes of
polyatomic alcohols in the liquid state. Zhur.fiz.khim.
37 no.2:347-353 F '63. (MIRA 16'5)

1. Moskovskiy gosudarstvennyy universitet imeni Lomonosova.
(Alcohols) (Dielectric constants)

KOMANDIN, A.V.; SIZOV, L.I.; SHIMIT, B.D. (Moscow)

Dielectric constant and dielectric losses of o-hydroxybenzoic
acid derivatives in the liquid state. Zhur. fiz. khim. 37
no.4:764-769 Ap '63. (MIRA 17:7)

1. Moskovskiy gosudarstvennyy universitet imeni M.V. Lomonosova.

SPS(a)/SPS(1)/SPS(2)/SPS(3)/SPS(4)/SPS(5)/SPS(6)/SPS(7)/SPS(8)/SPS(9)/SPS(10)/SPS(11)/SPS(12)/SPS(13)/SPS(14)/SPS(15)/SPS(16)/SPS(17)/SPS(18)/SPS(19)/SPS(20)/SPS(21)/SPS(22)/SPS(23)/SPS(24)/SPS(25)/SPS(26)/SPS(27)/SPS(28)/SPS(29)/SPS(30)/SPS(31)/SPS(32)/SPS(33)/SPS(34)/SPS(35)/SPS(36)/SPS(37)/SPS(38)/SPS(39)/SPS(40)/SPS(41)/SPS(42)/SPS(43)/SPS(44)/SPS(45)/SPS(46)/SPS(47)/SPS(48)/SPS(49)/SPS(50)/SPS(51)/SPS(52)/SPS(53)/SPS(54)/SPS(55)/SPS(56)/SPS(57)/SPS(58)/SPS(59)/SPS(60)/SPS(61)/SPS(62)/SPS(63)/SPS(64)/SPS(65)/SPS(66)/SPS(67)/SPS(68)/SPS(69)/SPS(70)/SPS(71)/SPS(72)/SPS(73)/SPS(74)/SPS(75)/SPS(76)/SPS(77)/SPS(78)/SPS(79)/SPS(80)/SPS(81)/SPS(82)/SPS(83)/SPS(84)/SPS(85)/SPS(86)/SPS(87)/SPS(88)/SPS(89)/SPS(90)/SPS(91)/SPS(92)/SPS(93)/SPS(94)/SPS(95)/SPS(96)/SPS(97)/SPS(98)/SPS(99)/SPS(100)

AUTHOR: Komandin, A. V.; Sizov, L. I.; Shimit, B. D.

76
75

TITLE: Thermodynamics of dielectric relaxation processes in liquids.

SOURCE: AN SSSR. Zhurnal fizicheskoy khimii v. 37 no. 5 1963, 1083-1088

TOPIC TAGS: relaxation processes phenyl O-hydroxybenzoate phenyl
O-ethoxybenzoate

SUMMARY: A previous study (A. V. Komandin, L. I. Sizov and B. D. Shimit, Zh. fizicheskoy khimii, 37, 264, 1963) was made on the temperature dependence of the dielectric relaxation of phenyl O-hydroxybenzoate and phenyl O-ethoxybenzoate at several temperatures. The present work is concerned with the study of the dielectric relaxations of these compounds at several temperatures. From the results of the present investigations, the main thermodynamic functions characterizing the dielectric relaxation processes in these compounds have been determined. The activation energy of the dielectric relaxation of phenyl O-ethoxybenzoate is 10.5 kcal/mole.

Card 1/2

L 9903-63

ACCESSION NR: AP3000418

of phenyl o-acetoxybenzoate at 55 and 42C in the supercooled state has been
determined. A linear relationship between $\log \tau$ and $1/T$ has been established
for these and their corresponding isomeric compounds. A detailed
analysis of the data is given. The results are presented in the form of
equations, 7 tables and 3 graphs.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet (Moscow State University)

SUBMITTED: 07May62 DATE ACQ: 19Jun63

ENCL: 00

SUB CODE: 00

NR REF SOV: 005

OTHER: 002

Card

2/2

KOMANDIN, A.V.; SHIMIT, B.D.

Thermodynamics of dielectric relaxation processes of o-hydroxybenzoic acid derivatives. Zhur.fiz.khim. 37 no.10:2289-2293. 0 '63.

(MIRA 17:2)

1. Moskovskiy gosudarstvennyy universitet.

ACCESSION NR: AP4011441

S/0076/64/038/001/0089/0095

AUTHORS: Komandin, A. V. (Moscow); Smirnova, A. D. (Moscow)

TITLE: The thermodynamic aspect of dielectric relaxation processes in glycerin esters.

SOURCE: Zhurnal fiz. khim, v. 38, no. 1, 1964, 89-95

TOPIC TAGS: dielectric constant, dielectric losses, glyceryl acetate, hydroxybenzoate, enthalpy, entropy, dielectric relaxation, acetic acid, glycerin derivatives, thermodynamic functions

ABSTRACT: This article deals with the measurements of the dielectric constants and tangent of dielectric losses of α, γ -glyceryl acetate and α -glycerin ester of o-hydrobenzoate in a liquid and supercooled liquid state at different frequencies of the electric field and in a wide range of temperatures. The free energy, enthalpy and entropy characterizing the dielectric relaxation processes were calculated from the resulting measurements. The relaxation time of the glycerin derivatives is closely associated with the nature of the substituting groups. The substitution of the hydrogen in the glycerin by simpler

Card 1/2

ACCESSION NR: AP4011441

radicals reduces the relaxation time, whereas the substitution by more complex radicals prolongs the dielectric relaxation time to a considerable extent. The dispersion of the dielectric constant of α, β -glyceryl acetate has been determined for five temperatures ranging from -20 to -50C, and the dispersion of α -glyceryl o -hydroxybenzoate has been determined at 40, 30 and 20C. The temperature dependence of the dielectric constant and the loss tangent have been determined by α -glyceryl o -hydroxybenzoate over a given temperature range. The connection between the thermodynamic functions and the chemical structure of the investigated compounds is under discussion. Orig. art. has: 3 Figures, 6 Formulas and 6 Tables.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet imeni V. M. Lomonosov (The V. M. Lomonosov Moscow state university)

SUBMITTED: 24Jan63

DATE ACQ: 14Feb64

ENCL: 00

SUB CODE: CH

NR REF SOV: 006

OTHER: 003

Card 2/2

ACCESSION NR: AF4033413

8/0076/64/038/003/0783/0785

AUTHOR: Komandin, A. V.; Smirnova, A. D.

TITLE: Dielectric properties of liquid benzophenone

SOURCE: Zhurnal fizicheskoy khimii, v. 38, no. 3, 1964, 783-785

TOPIC TAGS: benzophenone, dielectric loss, dielectric permeability, relaxation process, thermodynamic function

ABSTRACT: The article describes the measurements of dielectric permeability and the tangent of the angle of dielectric losses for liquid benzophenone in the 95 to -55 C interval at 50 kc, 800 kc and 20 Mc as the frequency of the external field. The measurement of dielectric properties was conducted by the resonance method. The accuracy of dielectric permeability measurement was $\pm 1\%$, dielectric losses $\pm(5 - 10)\%$, and the accuracy of temperature measurements was ± 0.1 deg. It was found that with lowering of the temperature the dielectric permeability of liquid benzophenone increases, reaching a maximum, after which it rapidly falls to small values, which characterizes the deformation polarization. At the temperature of the maximum of dielectric permeability the tangent of the dielectric

Card 1/3

ACCESSION NR: AP4033413

SUBMITTED: 16 May 63

SUB CODE: EM

NO REF SOV: 004

ENCL: 00


OTHER: 000

Card

3/3

"APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000824020006-9



APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000824020006-9"

Komandin, N. L.

137-1957-12-23429

Translation from: Referativnyy zhurnal, Metallurgiya, 1957, Nr 12, p 86 (USSR)

AUTHOR: Komandin, N. L.

TITLE: The Continuous Casting of Steel (Nepreryvnaya razlivka stali)

PERIODICAL: V sb.: Novoye v liteyn. proiz-ve. Nr 2. Gor'kiy, Knigoizdat, 1957, pp 150-171

ABSTRACT: An analysis of the operation of an industrial system, installed at the "Krasnoye Sormovo" plant in 1955, by means of which two ingots of 175 - 420 mm cross section can be cast simultaneously from a 45-50 t ladle through an intermediate casting device (ICD). These unified ingots replace 12 types of regular ingots of different dimensions and weights and are suitable for rolling into sheets and structural shapes. To ensure the completeness of the process of casting steel from a ladle of such capacity, the temperature of the metal must be 1540° or greater (as read on an optical pyrometer) at the time of its discharge from the furnace. If the temperature is lower, difficulties in the casting process are encountered because the metal solidifies in the casting nozzle of the ICD. An ICD of 3-5 t capacity, equipped with a cover and heated to a temperature of 900°, ensures the pouring of the

Card 1/3

137-1957-12-23429

The Continuous Casting of Steel

entire contents of the ladle at an almost constant temperature (the temperature drop in the ICD is only 8-10° in 45-60 min). In order to achieve a steady stream of metal from the ICD various casting nozzles were tested, including the electrically heated graphite-chamotte type. Compared with the regular ingot produced at the "Krasnoye Sormovo", the continuous ingot shows a finer grain structure, decreased liquation and porousness, and a reduced amount of non-metallic inclusions. However, some internal cracks were observed in it; the occurrence of the cracks depends on the intensity of the secondary cooling. A diagram is shown for the stresses which arise in the cross section of a continuous ingot. When the amount of water applied to the surface of the ingot in jets during the secondary cooling is decreased from 5 to 3.5 liter per kg of steel, the size of the cracks is considerably reduced. Internal cracks may be eliminated entirely if the surface of the ingot is sprayed by water from spray nozzles. The internal cracks are completely welded during the rolling of the ingots into sheets, and the surface of the sheets is superior to that of the sheets obtained from regular ingots. The mechanical properties of the metal are better than the specification properties required for a given grade of steel. The plant employs

Card 2/3

137-1957-12-23429

The Continuous Casting of Steel

the metal obtained from rolling of continuous ingots for shipbuilding. The installation at the "Krasnoye Sormovo" casts 20-25 percent of the total steel smelted in the Martin shop, and it is planned to increase this amount to 60-65 percent in the near future.

N. N.

1. Steel castings-Test methods
2. Steel castings-Test results

Card 3/3

137-58-4-8335

Translation from: Referativnyy zhurnal, Metallurgiya, 1958, Nr 4, p 289 (USSR)

AUTHORS: Komandin, N. L., Smolyakov, B. N.

TITLE: Shipbuilding Applications of High Strength Steels (Primeneniye staley povyshennoy prochnosti v sudostroyenii)

PERIODICAL: Tr. Gor'kovsk. in-ta inzh. vodn. transp., 1957, Nr 14, pp 62-75

ABSTRACT: The properties of MK steel of the % composition C 0.09-0.12, Si 0.7-1.2, Mn 1.0-1.6, and Cu 0.2-0.5 were investigated relative to its use in shipbuilding in the place of SKhL-1 and St. 3 steels. Testing of >2000 specimens cut from plates of different thicknesses along and across the direction of rolling have shown MK steel to have superior plastic properties than St. 3 steel, lower notch and aging sensitivity, better corrosion strength in sea and river water and in a mixture of 20% benzene and 80% sea water, and superior weldability. The building of a seagoing ship of MK steel saved 26.5% in weight; a 17.6% weight saving was attained in the building of a river vessel. V. L.

Card 1/1

1. Ship hulls--Materials--Effectiveness 2. Steel--Mechanical properties 3. Steel--Applications

Komandin, N.L.


KOMANDIN, N.L., kand.tekhn.nauk.

Use of SKhL-1 steel in shipbuilding. Rech.transp. 16 no.12:17-19
D '57. (MIRA 11:1)

(Shipbuilding--Supplies)
(Steel, Structural)

"APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000824020006-9



APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000824020006-9"

"APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000824020006-9

APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000824020006-9"

ZINEVICH, I.N.; KOMANDIROV, G.L.

Equipment for burnishing external cylindrical surfaces. Mashinostroitel'
no.4:2-3 Ap '65. (MIRA 18:5)

KOMANDIROV, Yu. (Khabarovsk)

Loudspeaker instead of microphone. Radio no. 7:34 J1 '54.
(MIRA 7:7)

(Loud-speakers) (Microphone)

Komandirov, Yu.

107-12-19/46

AUTHOR: Komandirov, Yu., Kemerovo

TITLE: Long-Distance Communications on the Ultrashort Waves
(Dal'niye svyazi na UKV)

PERIODICAL: Radio, 1956, Nr12, p. 20 (USSR)

ABSTRACT: A report of unusually good transmission of ultrashort waves on September 28 and 30, 1956, in the central part of Russia is presented.

Sept 28, 1956, the collective radio station of the DOSAAF Kemerovo oblast Radio Club 059510 established (at 11.52 Moscow time) a duplex communication with the station 068003 in Rostov-na-Donu, on 38 mc, at a distance of 3.200 km. The 068003 worked telegraphically with RST 338, and 056510 worked telephonically with RSM 595.

Sept 30, 1956, a duplex radio communication was established with Novocherkassk, 068030, 3120 km (at 08.43 a.m.).

At 09.00 a.m. the communication was established with Moscow 077528 2960 km. Moscow operator Ye. Skorospelov reported his RSM 445, and mine RSM 595.

At 09.40 a.m. the same day, a contact was established with the Taganrog station 068040, distance 3820 km, with RSM 575 and Taganrog RSM 445.

Card 1/2 Between 10.00 and 11.00 a.m. the same day, radio hams of Kemerovo heard,

107-12-19/46

Long-Distance Communications on the Ultrashort Waves

with RSM 335-575, the following stations: Ivanovo 057011, Saratov 069004 and 069007, Voroshilov 031482, Leningrad 076524, Silishchi 007001, Khimki 064020, and many other cities.

The DOSAAF Club Kemerovo station has a 9-tube superhet receiver.

AVAILABLE: Library of Congress

Card 2/2

8/138/60/000/01/01/010

AUTHORS: Shatalov, V.A., Popova, Ye.N., Krygina, K.G., Komandorova, L.A.

TITLE: Greater Stability of the SKS-30 Polymerization System

PERIODICAL: Kauchuk i Rezina, 1960, ¹⁹No. 1, pp. 3 - 5

TEXT: The article deals with possibilities of improving the stability of the polymerization system and investigates the reasons for its instability. One of the main reasons for the separation of coagulum during the process of polymerization is due to ferrous salt getting into the polymerization charge. It is sufficient to add 0.1% FeSO₄ to the aqueous phase to bring the amount of coagulum from 0.5 to 4.7% based on the rubber. The principal mass of iron compounds enters the polymerization mixture with the emulsifier (Nekal) and alkali. Deposition of coagulum is also brought about by a great temperature gradient between the latex and the walls of the apparatus. The large particles of phenyl- β -naphthylamine are the coagulation centers. The stability of the low-temperature SKS-30A polymerization system can be improved by increasing the quantity of water and emulsifier, as well as by adding a small amount of leucanol, which should be introduced at a rate of 0.3% based on the monomers. The effect produced by leucanol on the SKS-30 system

Card 1/2

Greater Stability of the SKS-30 Polymerization System

S/138/60/000/01/01/010

has been tested under laboratory and industrial conditions. The results of tests are shown in 2 Tables. These data show that under industrial conditions leucanol improved considerably the stability of the polymerization system. After introduction of leucanol the use of the deposition in the end polymerizers decreased about 10 times, while in the first apparatus coagulation was practically not existing. Laboratory tests permitted to draw the conclusion, that the stabilization brought about by leucanol is due to the effect it produces on the ion of iron and to the physico-chemical processes of colloidal substances like soap or dispersers, where- by the protective action of the film surrounding the rubber particles is strengthened. It can therefore be concluded that by the introduction of leucanol into the recipe of SKS-30, by the total prevention of iron compounds from getting into the system and by the improvement of the dispersion of phenyl- β -naphtylamine it is possible to eliminate the precipitation of coagulum from latex in the course of polymerization as well as the separation of monomers. There are 3 tables and 3 Soviet references. ✓

ASSOCIATION: Voronezhskiy zavod sinteticheskogo kauchuka im. S. M. Kirova
(Voronezh Plant of Synthetic Rubber im. S. M. Kirov)

Card 2/2

KOMANDRINA, T.A. [Komandrina, T.O.]

New method for calculating seismic effects on spatial constructions.
(MIRA 17:9)
Dop. AN URSR no.7:888-892 '64.

1. Odesskiy inzhenerno-stroitel'nyy institut. Predstavleno
akademikom AN UkrSSR G.N.Savinym [Savin, H.M.].

YEGUPOV, Vyacheslav Konstantinovich. Prinimala uchastiye
KOMANDRINA, T.A.; KIYANICHENKO, N.S., red.

[Calculating buildings for strength, rigidity, and vibration] Raschet zdanii na prochnost', ustoychivost' i kolebaniia. Kiev, Budivel'nyk, 1965. 253 p. (MIRA 18:7)

KOMANDRINA, T.A.

Calculating buildings for seismic action taking three-dimensional effect into account. Izv. AN Arm. SSR. Ser. tekhn. nauk 17 no.4:17-25 '64. (MIRA 17:11)

1. Odesskiy inzhenerno-stroitel'nyy institut.

^{REV}
KOMANDERSKAYA, L. V.

USSR/ Physical Chemistry - Thermodynamics. Thermochemistry. Equilibrium.
Physicochemical analysis. Phase transitions

B-8

Abs Jour : Referat Zhur - Khimiya, No 4, 1957, 11143

Author : 1. Shukarev S.A., Tolmacheva T.A., Oranskaya M.A., Komanderskaya L.V.
2. Shchukarec S.A., Oranskaya M.A., Shemyakina T.C.

Title : Thermal Dissociation of Platinum Hakides. Communication 1. Platinum
Bromides. Communication 2. Platinum Chlorides.

Orig Pub : Zh. neorgan. khimii, 1956, 1, No 1, 8-16; 17-23

Abstract : 1. Statistical method of F. Ephraim (Ber., 1917, 1069) was used to investigate temperature dependence of thermal dissociation of $PtBr_4$ (I), $PtBr_3$ (II), $PtBr_2$ (III) and $PtBr$ (IV). Scheme of the unit is described. Data obtained are represented as $\lg P - 1/T$ graphs. From the slope of the straight lines were determined equations of dependence of dissociation pressure on temperature, for I $\lg P = 7.809 - (4549/T)$, II $\lg P = 7.195 - (4808/T)$, III $\lg P = 6.064 - (5123/T)$ and IV $\lg P = 4.755 - (4679/T)$. Calculated therefrom were the values of P_{Br_2} at 10° intervals, for I (over range 200-280°), II (280-390°), III (420-500° and IV (460-510°) which are tabulated. By using these data calculations were made of the values of

Card 1/3

USSR/ Physical Chemistry - Thermodynamics. Thermochemistry. Equilibrium.
Physicochemical analysis. Phase transitions

B-8

Abs Jour : Referat Zhur - Khimiya, No 4, 1957, 11143

enthalpy and entropy changes (ΔH and ΔS) in reactions of successive dissociation of the bromides, on the assumption that these quantities are independent of the temperature within the interval under study. Determined were the values of enthalpy of formation ΔH (formation), (in kcal/mole) from metal and gaseous bromine, for I, II, III and IV, which are, respectively, -44.0; -33.6; -22.6; -10.75, and thus in good agreement with literature data. The actual existence of each bromide was confirmed by chemical analysis and recording of Debye X-ray patterns. It is shown that below 340°K IV must undergo exothermal disproportionation. The appreciable scattering of experimental data forces the authors to assume that platinum bromides possess the property of interacting with one another to form solid solutions. 2. Investigated was the temperature dependence of dissociation pressure of $PtCl_4$ (V) (298-358°), $PtCl_3$ (VI) (332-394°), $PtCl_2$ (VII) (490-530°), and $PtCl$ (VIII) (568-762°). On the basis of the data thus obtained the enthalpies of formation were calculated (kcal/mole): V -62.7; VI -48.1; VII -33.4; VIII -11.7; these values are in satisfactory agreement with literature data. Debye X-ray patterns of the platinum chlorides have

Card 2/3

APPROVED FOR RELEASE: 06/13/2000 CIA-RDP86-00513R000824020006-9"
USSR/ Physical Chemistry - Thermodynamics. Thermochemistry. Equilibrium.
Physicochemical analysis. Phase transitions

B-8

Abs Jour : Referat Zhur - Khimiya, No 4, 1957, 11143

demonstrated the actual existence of each substance. Experimental data for the reactions $2PtCl_n = 2PtCl_{n-1} + Cl_2$ are expressed by the equations $\lg P_{Cl_2} = A - (B/T)$, wherein A and B are: for $n=4$, 9.04, 6267.5; $n=3$, 8.42, 6303.1; $n=2$, 10.59, 9131.1; $n=1$, 4.36, 5114.9. On the basis of the data obtained and those found on the literature an analysis is made of the dependence of isobaric potential of the valency is discussed. It is shown that at low temperatures ($< 600^\circ$) VIII must undergo exothermal disproportionation.

Card 3/3

KOMANDROVSKIY, V. G.

PHASE I BOOK EXPLOITATION

807/6100

Akademiya nauk SSSR. Institut tochnoy mekhaniki i vychislitel'noy tekhniki.

Trudy (Academy of Sciences of the USSR, Institute of Precision Mechanics and Computer Technology. Transactions) no. 2. Moscow, 1961. 447 p. 1000 copies printed. Contributors not mentioned.

PURPOSE: This collection of articles is intended for scientific and technical personnel concerned with machine translation and computer technology.

COVERAGE: This collection of articles of the Institute of Precision Mechanics and Computer Technology, Academy of Sciences USSR, is the second in a series concerned with machine translation and mathematical linguistics. The collection contains reports written by members of the Machine-Translation Group of the Institute as well as reports by researchers from other organizations. The articles deal with various problems in machine translation, such as the possibility of an intermediate language, relationships between various languages, systems of recording, structure of

Card 1/6

Card 6/6

IS, wrc/eb
11/30/62

Academy of Sciences (Cont.)

SOV/6100

26. Ivanov, V. V. On the Acceptability of Phonological Patterns 398

27. Yefimov, M. B., and A. A. Zvonov. Attempt at Constructing a System of Graphic Analysis of Hieroglyphic Writing 415

28. Komandrovskiy, V. G. Problems of Constructing Reading Device 425

References

444

AVAILABLE: Library of Congress

SUBJECT: Automation and Computer Engineering

Card 6/6

IS/wrc/eb
11/30/62

L 33746-66 EWT(d)/T/EWP(1) IJP(c) BB/GG/UD/JAT(BF)
ACC NR: AT6008567 SOURCE CODE: UR/0000/65/000/000/0169/0178

AUTHOR: Komandrovskiy, V. G.

52
Bx1

ORG: none

TITLE: A logical scheme for symbol identification 16C

SOURCE: AN SSSR. Institut nauchnoy informatsii. Chitayushchiye ustroystva (Reading devices). Moscow, VINITI, 1965, 169-178

TOPIC TAGS: pattern recognition, character reading equipment, reading machine,
STATISTICAL DISTRIBUTION

ABSTRACT: A logical scheme for identifying printed symbols on the basis of statistical features is presented. The scheme eliminates certain problems arising from varying size, slant and other differences in printed symbols. The method considers text symbols from the aspect of their length, width of the contour, minimum number of scanning lines needed for recognition, slope of the line, and other symbol properties. These properties are translated into statistical distributions of black-white fields. The final step of the method involves comparison with some standard introduced into the machine. An apriori determination of the statistical probability of distribution introduces an elementary self-reading aspect into the recognition process. An example is given in which an entire handwritten word is analyzed and its parameters are given in tabular form. Orig. art. has: 7 figures, 2 tables.

SUB CODE: 00/ SUM DATE: 09Sep65/ ORIG REF: 001/ OTH REF: 003

Card 1/1 PLG

TRCA, S.; MISINGER, J.; KOMANEC, J.

Apropos of the relation between the pathogenesis of inflammation after induced abortion and the duration of pregnancy. Cesk. gynek. 29 no.8:613-616 0 '64.

1. II. gyn.-por. klin. fak. vseob. lek. Karlovy University v Praze, (prednosta prof. dr. J. Lukas, DrSc.); Ustr. ustav zdravotnicke osvety v Praze, (reditelka MUDr. M. Taufrova, CSc.).

DOLEZAL, A., CSc.; KOMANEČ, J.; TRNKA, V., doc, CSc.

Some considerations on causes of failure in the treatment of threat to the pregnancy. Cesk. gyn. 27 [41] no.6/7:439-445 Ag '62.

1. II. gyn.-por. klin. fak. vseob. lek. KU v Praze, prodnosta prof.
dr. J. Lukas, DrSc.

(ABORTION THREATENED)

JAROS, Emil, inz.; KOMANEK, Zdenek, inz.

Technological projects of specialized centers. Stroj vyr
12 no.3:202 '64.

1. Branch of the Czechoslovak Scientific Technological Society
at the Zavody presneho strojirenstvi, National Enterprise,
Gottwaldov.

GONIKBERG, M.G.; DOROGCHINSKIY, A.Z.; GAVRILOVA, A.Ye.; KOMANENKOVA, R.A.;
MITROFANOV, M.G.; KUPRIYANOV, V.A.

Determination of the naphthalene and alkyl naphthalene content of
stocks and dealkylation products. Neftekhimiia 3 no.6:916-921 N-D
'63. (MIRA 17:3)

1. Institut organicheskoy khimii AN SSSR im. N.D.Zelinskogo i
Groznskiy neftyanoy nauchno-issledovatel'skiy institut.

GONIKBERG, M.G.; GAVRILOVA, A. Ye.; ALEKSEYEV, Ye.F.; KOMANENKOVA, R.A.

Homogenous demethylation of methyl naphthenes. Neftekhimiya
4 no.2: 252-256 Mr-Ap'64 (MIRA 17:8)

1. Institut organicheskoy khimii AN SSSR imeni Zelinskogo

Komanich, D.


APPROVED FOR RELEASE: 06/13/2000 v. CIA-RDP86-00513R000824020006-9"

AUTHORS: Petrov, A. D., Mironov, V. P.,
Komanich, D.

TITLE: Synthesis of β -Chloro and β , β -Dichlorovinylalkyldichlorosilanes (Sintez β -Khlor- 1 β , β -dikhlorvinilalkildikhlorosilanov).

PERIODICAL: Izvestiya AN SSSR, Otdelenie Khimicheskikh Nauk, 1957,
Nr 11, pp. 1393-1395 (USSR)

ABSTRACT: Here the best conditions for a synthesis of $\text{ClCH} = \text{CHSiCl}_2$, as well as the alkyl- and alkoxy-derivatives of them, among them also the difunctional (in the silicon atom) β -chlorovinylethylenedichlorosilane, were obtained. The development of C. L. Agre's (reference 2) was continued here and it was ascertained that the trichloroethylene can also be condensed with the alkyldichlorosilanes. This is a very simple way for the synthesis of the dichlorodialkylsilanes with the functional groups in the alkyl-chain



AGARBICHANU, I.I. [Agarbiceanu, I.]; BLANARU, L.; DRAGANESKU, V.;
IONESCU-PALLAS, N.I. [Ionesco-Pallas, N.J.]; KOMANICHU, N.;
TATU, V.

Determining the nuclear magnetic moment of the isotope Hg^{199} from
the hyperfine structure of the HgI 5461 Å line. Opt. i spektr. 10
no.3:297-300 Mr '61. (MIRA 14:8)

1. Institut atomnoy fiziki AN Rumynskoy Narodnoy Respubliki,
Bukharest.

(Nuclear moments) (Mercury--Isotopes) (Interferometry)

ACCESSION NR: AP4020919

S/0051/64/016/002/0182/0186

AUTHOR: Draganesku, V.; Komanichu, N.; Tatu, V.; Ionescu-Pallas, N.Zh.

TITLE: Hyperfine and isotope structure of the 5535 Angstrom line of Ba I

SOURCE: Optika i spektroskopiya, v.16, no.2, 1964, 182-186

TOPIC TAGS: hyperfine structure, isotope structure, isotope shift, barium(I), barium(I) resonance line, hfs theory, LS coupling calculations

ABSTRACT: The 5535 Å Ba I line has been investigated by H.Kopefermann and O.Wessel (Nachr.Akad.Wissensch.,Göttingen,Math.-Phys.No.2,53,1948), O.H.Arroe (Phys.Rev.,79, 836,1950) and later by D.A.Jackson (Ibid.,106,948,1957 and Proc.Roy.Soc.,A263,289, 1961), but only the last, who used absorption in an atomic beam, was able to discern hyperfine structure. The present theoretical and experimental investigation of this resonance line, associated with the $Gs^2 1S_0 - Gs6p^1P_1$ transition, was undertaken to check the experimental data and provide a theoretical explanation thereof. (Jackson noted a discrepancy between the observed and calculated intensities and hypothesized a small hfs (10 mK) for the odd isotopes.) The present experimental work was also carried out by the absorption in an atomic beam technique; the setup

Card 1/3

ACCESSION NR: APL020919

for obtaining the atomic beam was an improved version of the apparatus described by V. Draganescu (Opt. i spektro., 10, 301, 1961) and included provision for water-cooling of the evaporator in order to enhance the vacuum. A special device consisting of two mirrors with slits was employed to make the light beam pass several times through the atomic beam, in order to obtain substantial absorption without recourse to increasing the evaporator temperature. The light source was a hollow cathode tube into which the barium was introduced in the form of BaCl_2 . The spectroscopic instrument was an ISP-22 spectrograph crossed with a Fabry-Perot etalon; the interference patterns were photographed with an exposure of under 15 min and scanned on a Zeiss microphotometer. There were resolved in all 7 components arranged almost symmetrically with respect to the central 138 component in an interval of about 40 mK ($1 \text{ mK} = 10^{-3} \text{ cm}^{-1}$). The experimental data are compared in a table with the results of calculations based on the intermediate coupling theory of G. Breit and L. A. Wills (Phys. Rev. 44, 470, 1933), assuming Russell-Saunders (L-S) coupling. The computation formulas are adduced, and corrections are made for finite nuclear size and the nuclear magnetism distribution. The agreement between the experimental data and calculated values is satisfactory. The theory at present is not adequate for deciding for or against Jackson's hypothesis. The proposed interpretation allows of explaining some

Car 2/3

ACCESSION NR: APL020919

of the observed components. Nevertheless further measurements aimed at determining the broadening of the hfs in the spectra of separate isotopes are planned. "The authors are grateful to I. Agarbichan for useful discussions during the course of the work." Orig.art.has: 17 formulas and 1 table.

ASSOCIATION: Institut atomnei fiziki, Bucharest (Institute of Atomic Physics)

SUBMITTED: 10May63

DATE ACQ: 02Apr64

ENCL: 00

SUB CODE: PH

NR REF SOV: 003

OTHER: 009

3/3
Card

KOMANITSKY, V.; PLATZEROVA, L.

Determining of the fractional composition of titanium dioxide.
Khim. volok. no.6:38-41 '65. (MIRA 18:12)

1. Narodnoye predpriyatiye KhEMKO, zavod Gummene, Chekhoslo-
vatskaya Sotsialisticheskaya Respublika.

KOMANOV, A. (Major)

"Problems of Utilization of Atomic Energy," Part I., Krasnaya Zvezda, No.8,
page 3, January 11, 1955

Summary of an article - D 230702, 12 May 55

ROMANOV, B.O.; SHAKIROV, E.Sh.

Mesozoic age of magnetites from the skarn formations of the
Cava deposit. Dokl. AN SSSR 160 no.6:1378-1380 F '65.

(MIRA 18:2)

1. Submitted June 30, 1964.